



Technological University Dublin
ARROW@TU Dublin

Articles

NanoLab

2016-06-01

Modification of the In Vitro Uptake Mechanism and Anti-Oxidant Levels in HaCaT Cells and Resultant Changes to Toxicity and Oxidative Stress of G4 and G6 Poly (amido amine) Dendrimer Nanoparticles.

Marcus Maher

Technological University Dublin

Hugh J. Byrne

Technological University Dublin, hugh.byrne@tudublin.ie

Follow this and additional works at: <https://arrow.tudublin.ie/nanolart>



Part of the [Other Biochemistry, Biophysics, and Structural Biology Commons](#), and the [Toxicology Commons](#)

Recommended Citation

Maher, M. and Byrne, H., (2016) Modification of the in vitro uptake mechanism and anti-oxidant levels in HaCaT cells and resultant changes to toxicity and oxidative stress of G4 and G6 Poly (amido amine) dendrimer nanoparticles. *Analytical and Bioanalytical Chemistry*, 408, 5295-5307 (2016)

This Article is brought to you for free and open access by the NanoLab at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact yvonne.desmond@tudublin.ie, arrow.admin@tudublin.ie, brian.widdis@tudublin.ie.



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#)



Modification of the *in vitro* uptake mechanism and anti-oxidant levels in HaCaT cells and resultant changes to toxicity and oxidative stress of G4 and G6 Poly (amido amine) dendrimer nanoparticles.

Marcus A. Maher^{1,2,*}, Hugh J. Byrne¹

¹FOCAS Research Institute, Dublin Institute of Technology, Kevin Street, Dublin 8, Ireland

²School of Physics, Dublin Institute of Technology, Kevin Street, Dublin 8, Ireland

*Corresponding author: marcus.maher@mydit.ie

Keywords:

Dendrimer nanoparticles

PAMAM nanoparticles

Endocytosis

Passive diffusion

Oxidative stress

Membrane permeability

Abstract:

The mechanism of cellular uptake by endocytosis and subsequent oxidative stress has been identified as the paradigm for the toxic response of cationically surface charged nanoparticles. In an attempt to circumvent the process, the effect of increased cellular membrane permeability on the uptake mechanisms of poly (amidoamine) dendrimers generation 4 (G4) and 6 (G6) *in vitro* was investigated. Immortalised, non-cancerous human keratinocyte (HaCaT) cells were treated with DL-Buthionine-(S,R)-sulfoximine (BSO). Active uptake of the particles was monitored using fluorescence microscopy to identify and quantify endosomal activity and resultant oxidative stress, manifest as increased levels of reactive oxygen species, monitored using the carboxy-H₂DCFDA dye. Dose dependent cytotoxicity for G4 and G6 exposure was registered using the cytotoxicity assays Alamar Blue and MTT, from 6 to 72 hours.

Reduced uptake by endocytosis is observed for both dendrimer species. A dramatic change, compared to untreated cells, is observed in the cytotoxic and oxidative stress response of the BSO treated cells. The significantly increased mitochondrial activity, dose dependent anti-oxidant behaviour and reduced degree of endocytosis for both dendrimer generations, in BSO treated cells, indicates enhanced permeability of the cell membrane, resulting in the passive, diffusive uptake of dendrimers, replacing endocytosis as the primary uptake mechanism. The complex MTT response reflects the importance of glutathione in maintaining redox balance within the mitochondria. The study highlights the importance of regulation of this redox balance for cell metabolism, but also points to the potential of controlling the nanoparticle uptake mechanisms, and resultant cytotoxicity, with implications for nanomedicine.

Introduction

Nanoparticle science is a rapidly advancing field which holds much promise in areas such as targeted drug and gene delivery.^{1,2,3} However, nanoparticle uptake into a cell has been demonstrated, particularly for nanoparticles with an effective cationic surface charge, to give rise to cytotoxic responses, raising concerns about the potential health and environmental impact of the proliferation of nanomaterials in consumer products.⁴ A systematic understanding of the mechanisms of toxicity and their dependence on nanoparticle physico-chemical properties on a cellular level is therefore required.⁵ In the context of nanomedical applications, understanding and controlling the uptake process and subcellular trafficking of the delivery vehicle and the bioavailability of the cargo are critically important.

Cellular uptake of nanoparticles principally occurs via endocytosis, whereby the nanoparticle is invaginated by the cellular membrane and is transported into the cell.⁶ As the low pH environment of the endosome attempts to digest the nanoparticle, the redox balance of the cell is disrupted, and, in the case of nanoparticles with an effective cationic surface charge, the process gives rise to an increase in the production of Reactive Oxygen Species (ROS), localised mainly around the endosome (or later lysosome).^{6,7,8,9} Although intra cellular anti-oxidants attempt to neutralise the imbalance, ROS production can be sufficient to lead to oxidative stress.¹⁰ Subsequently, a cascade of events and the release of several characteristic cytokines and chemokines occurs, ultimately leading to cell death, a process which is the accepted paradigm of nanotoxicity of many nanoparticles *in vitro*.⁶ The process has been well demonstrated for model nanoparticle system such as amine functionalised polystyrene,^{11,12,13} amorphous nanosilica,¹⁴ and nanomeric polymeric dendrimers.¹⁰

Aminated molecules are intrinsic antioxidants, however, and generally regarded as ROS scavengers. As examples, spermine and spermidine have both been shown to reduce Fe^{3+} to Fe^{2+} ¹⁵ and the ferric reducing activity of these molecules has been identified as a measure of anti-oxidant potential.¹⁶ Carnosine, an endogenous dipeptide, has been shown to scavenge both reactive oxygen and nitrogen species.¹⁷ Notably, in a study by Khalid *et al.* (2015), while the larger, higher generations of the aminated nanoscale dendrimers Poly (propylene imine) (PPI) were demonstrated to elicit oxidative stress and significant toxicity, the smaller, lower generations exhibited intracellular antioxidant behaviour and low toxicity.¹⁸ Examination of the uptake mechanisms indicated a transition from cellular uptake by passive diffusion at low generations to active endocytosis for higher generations. The study suggests that, in the case of catatonically charge nanoparticles, the endocytotic uptake and trafficking process is, in itself, a source of cellular toxicity. In the context of drug delivery, invagination of the delivery vehicle and cargo in this harsh environment may also be undesirable, and although escape by endosomolysis is a potential strategy,¹⁹ this too can be a harsh process, causing significant damage to the cell.^{20,21} For these reasons, circumventing endocytosis appears to be a valid strategy for the reduction of toxicity associated with aminated nanoparticles as well as nanoparticle drug or gene delivery.

Guarnieri *et al.* (2015), demonstrated that functionalization of aminated polystyrene nanoparticles with the viral peptide gH625 (derived from *Herpes simplex virus – 1*), which has a membrane perturbing domain, enables translocation of particles to the cytoplasm, avoiding endocytosis and thus, dramatically reducing the cytotoxicity.²² An alternative strategy to increase

the permeability of the cellular membrane, *in vitro*, is through the application of DL-Buthionine-
(*S,R*)-sulfoximine (BSO). BSO has previously been employed to study the effects of the
reduction of levels of the intracellular antioxidant glutathione (GSH); and therefore oxidative
stress^{23,24,25,26} and has been shown to cause membrane permeabilisation.^{27,28} BSO works as an
inhibitor of the enzyme Glutamate Cysteine Ligase (EC 6.3.2.2) (historically known as gamma-
glutamylcysteine synthetase) which is the first step in the production of GSH in the cell.²⁹ This
reduction in GSH causes several different effects, one being the induction of the membrane
permeability transition in the mitochondria.^{30,31,32} The opening of this pore and the depletion of
GSH allows ROS to diffuse from the mitochondria to the cell and via lipid peroxidation, cause
damage to the cell membrane,²⁸ leading to increased permeability.

The reduction of GSH in the cell can have other adverse effects, however, mainly due to the fact
that GSH is one of the main antioxidants involved in maintaining the redox balance of the
mitochondria.³³ In the mitochondria, endogenous ROS is produced as a by-product of normal
metabolism and therefore completely eliminating GSH can leave the cell susceptible to damage
from this ROS.³⁴ Furthermore, reductions in cellular GSH can also lead to changes in the
regulation of Ca²⁺ distribution^{35,36,37} and the activation/deactivation of signalling pathways
involved with growth, differentiation and apoptosis.^{33,38,39,40} The loss of GSH and subsequent
changes in the cell have been implicated in several disease states which is reviewed elsewhere.³⁴

This study explores the impact, for HaCaT cell, of BSO treatment on the cellular uptake of, and
subsequent oxidative stress and toxic response to, poly (amido amine) (PAMAM) dendrimers.
These nanoscale aminated dendrimers have a systematically variable molecular structure, and the

111 homologous series of increasing generation, and concomitant size and number of surface amino
112 groups, has been demonstrated to be ideal to study the dependence of nanoparticle cellular
113 interactions on the physico-chemical properties of the particles. Previous studies have examined
114 the structurally dependent toxicity, and underlying mechanisms of endocytosis, oxidative stress,
115 immune responses and consequent toxicity^{7,8,9,10} and the responses have been numerically
116 modelled, as a guide towards a predictive toxicology approach.^{10,41} For consistency and to allow
117 comparisons with the previous work, the *in vitro* studies reported here were also carried out
118 using the immortalised human keratinocyte cell line, HaCaT, and identical oxidative stress and
119 cytotoxic assays. Furthermore, PAMAM dendrimers of Generation 4 and 6 were chosen as the
120 extremes of the previously reported cytotoxicological response. In comparison, it is
121 demonstrated that treatment with BSO results in a significant change in the nanoparticle uptake
122 mechanisms and cytotoxicity.

123 Materials and Methods

124 Materials

125 DMEM F12 HAM growth medium, Penicillin, Streptomycin, fluorescently labelled, Polystyrene
126 nanoparticles, with amine surface modification (PSNP-NH₂) 100nm, DL-Buthionine-(*S,R*)-
127 sulfoximine(BSO) and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) dye
128 were purchased from Sigma-Aldrich, Ireland. The PAMAM dendrimer nanoparticles,
129 generations 4 (molecular weight: 14,214 g/mol) and 6 (molecular weight: 58,046 g/mol), were
130 purchased from Sigma-Aldrich and manufactured by Dendritech Inc. ThiolTracker™ Violet
131 (TTV), CellLight® Early Endosomes-RFP, BacMam 2.0, Fetal Bovine Serum (FBS), L-
132 Glutamine, Alamar Blue (AB) and 5-(and-6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate
133 (carboxy-H₂DCFDA) dye were purchased from Life Technologies™, Bio-Sciences, Ireland.
134 HaCaT cells were purchased from Cell Line Services (CLS), Eppelheim, Germany. TrueLine 96-
135 well cell culture plates were used for all viability and ROS studies.
136 All fluorescence and absorbance readings were taken on a Molecular Devices SpectraMax M3
137 Spectrometer. Confocal Laser Scanning Fluorescence Microscopy (CLSM) images were taken
138 on a Zeiss LSM 510 Confocal Laser Scanning Microscope and processed using ImageJ software
139 (with co-localisation analysis performed with the JaCoP plugin for ImageJ). All viability, ROS
140 and GSH data analysis was performed using SigmaPlot v10.0 software.

141

142 Methods

143 Cell culture

144 HaCaT cells are an immortalised, non cancerous human keratinocyte cell line and were used for

these experiments. The cells were cultured in DMEM F12 HAM supplemented with 10% FBS, 45 IU/mL penicillin, 45 IU/mL streptomycin and 2mM L-glutamine at 37°C in 5% CO₂. All assays carried out in this set of experiments were performed in 96 well plates, with cells plated at a concentration of 1x10⁴ cells/well in 100μL of DMEM medium. Cells were allowed 24 hours to attach and were then treated with BSO for an additional 18 hours (at a concentration of 200μM), after which cells were exposed to PAMAM G4 or G6 dendrimers (in DMEM F12 HAM, with 5%FBS, 45 IU/mL penicillin, 45 IU/mL streptomycin, 2mM L-glutamine and 200μM BSO) at various concentrations for the specified time points. Six replicates of each concentration were performed per plate and each plate was repeated in triplicate.

ThiolTracker™ Violet (TTV)

ThiolTracker™ Violet (TTV) is a GSH detection agent. Cells were plated as described above and a concentration gradient of BSO was applied. Cells were left for 18 hours at 37°C in 5% CO₂ to allow for reduction of the amount of intracellular GSH. Cells were then washed twice with PBS, 100μL of TTV dye (at a final concentration of 20μM) were added to each well and the plates were allowed to incubate at 37°C in 5% CO₂ for 30 minutes, after which the TTV solution was removed and replaced with PBS. The fluorescence of each well was then read using the SpectraMax M3 spectrometer with λ_{EX}= 404nm and λ_{EM}= 526nm. GSH values were calculated as compared to the unexposed control.

Viability assays

Alamar Blue and MTT assays were used to determine the changes in cell viability, after treatment with BSO as described above, as a result of exposure to both PAMAM G4 and G6

dendrimers. Both Alamar Blue and MTT were performed on the same plate. The PAMAM G4 concentrations used were: 0.16, 0.32, 0.65, 1.3, 2.6, 5.2, 7.8 and 10.4 μ M, while the PAMAM G6 concentrations were: 0.08, 0.16, 0.32, 0.65, 1.3, 2.6, 3.9 and 5.2 μ M. The lower initial value of the PAMAM G6 dendrimers was used due to their reported EC₅₀ value being much lower than their G4 counterparts.^{7,8,9} Dose dependent viability percentages were calculated at time points: 6, 12, 24, 48 and 72 hours. Percentage viability was calculated as compared to a control which had been exposed to 200 μ M BSO, but had no nanoparticle treatment; this was to ensure any changes were caused by the nanoparticle and were not the result of BSO treatment. A separate control where no BSO was present was also performed and showed there was little difference between cells with no BSO exposure and cells which were exposed to BSO (Supplementary Material, Figure: S1).

Alamar Blue (AB)

The Alamar Blue assay was made up from 10X stock solution in medium (DMEM F12 HAM, with no additional supplements). At the specified time point, the plates were removed from the incubator and the medium containing PAMAM dendrimer was removed, the cells were washed with 100 μ L PBS and then 100 μ L of DMEM F12 HAM(unsupplemented) containing Alamar Blue were added to each well. The plates were incubated for 3 hours at 37°C in 5% CO₂ to allow for conversion of the dye. The fluorescence of each well was then read using the SpectraMax M3 spectrometer with λ_{EX} = 555nm and λ_{EM} = 585nm.

MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)

A stock solution of MTT was made at a concentration of 0.5mg/mL. 500 μ L of this stock were added for every 10mL of medium (DMEM F12 HAM, with no additional supplements). At the specified time point, the plates were removed from the incubator and the medium containing PAMAM dendrimer was discarded, the cells were washed with 100 μ L PBS and then 100 μ L of DMEM (unsupplemented) containing MTT was added to each well. The plates were incubated for 3 hours at 37°C in 5% CO₂ to allow for conversion of the dye. After 3 hours, any remaining dye was removed and each well was again washed with 100 μ L PBS, after which 100 μ L of DMSO was added and the plates were placed on a shaker for 10 minutes to allow for the dye to solubilise. The absorbance of each well was then read using the SpectraMax M3 spectrometer with $\lambda_{\text{ABS}}=595\text{nm}$.

Reactive Oxygen Species (ROS)

5-(and-6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate (carboxy-H₂DCFDA) dye was used for the detection of ROS. The dye was made up to a final concentration of 10 μ M in sterile PBS. Before addition of PAMAM dendrimer, this dye was added to the cells and allowed to incubate for 1 hour, after which, the dye was removed, the cells were washed thrice with PBS and the medium containing PAMAM dendrimer was added. At the specified time points the fluorescence was read by the SpectraMax M3 spectrometer with $\lambda_{\text{EX}}=488\text{nm}$ and $\lambda_{\text{EM}}=535\text{nm}$.

Confocal Laser Scanning Microscopy (CLSM)

Cells were plated onto MatTek 35mm glass bottom dishes at a concentration of 20,000 cells/dish in DMEM F12 HAM (supplemented with 10% FBS, 45 IU/mL penicillin, 45 IU/mL streptomycin and 2mM L-glutamine) and allowed to attach for 4 hours, at which point the

medium was removed and replaced with medium containing Celllight[®] Early Endosomes-RFP, BacMam 2.0 at a concentration of 20 particles per cell. Early endosome formation was tracked with the Celllight[®] Early Endosome – RFP kit, which transfects, into the cell, a version of Rab5a with a bound Red Fluorescent Protein. The cells were allowed to incubate (37°C, 5% CO₂) for 16 hours to ensure transfection with the early endosome reagent. After this, the medium was removed and cells were washed twice with PBS. For cells being tested without BSO, medium was added for 18 hours (these samples are referred to as untreated cells/untreated controls in the text), while for cells being tested with BSO, medium containing 200µM BSO was added for 18 hours. Cells were again washed with PBS and carboxy-H₂DCFDA was added for 1 hour (10µM in 2mL PBS), after which cells were again washed twice with PBS. PAMAM dendrimers were added at a concentration of 3.21µM (G4) and 1µM (G6) and cells were allowed to incubate for 3 hours (G4) or 1 hour (G6) and were then washed twice with PBS and imaged with the Zeiss LSM 510 Confocal Laser Scanning Microscope. 100nm PSNP-NH₂ with attached Green Florescent Protein was used as a positive control to ensure the Celllight[®] Early Endosomes-RFP was functioning as expected; the results of this test are available in the supplementary material section, Figure S2. Negative controls were also performed with cells which were not exposed to any nanoparticles, and little to no fluorescence was noted (data not shown). For ROS monitoring, doses and time points noted above were chosen based on the maximum responses previously reported in literature.⁸ All confocal images were analysed using ImageJ and co-localisation studies were performed using Manders split coefficients calculated with the JaCoP plugin for ImageJ.⁴²

Data Analysis and Statistics

236 Data analysis was performed using SigmaPlot™ v10.0 and fluorescence was calculated based on
237 the values of BSO controls (which were unexposed to nanoparticles, but had been treated with
238 200μM BSO for 18 hours).

239 “The cytotoxicity, GSH and ROS experiments were performed in 96-well microplates with six
240 replicates per plate and each plate repeated three times. Therefore, data points shown represent
241 the mean of 18 points, with error bars representing \pm the standard deviation (as calculated by
242 SigmaPlot™ v10.0). Confocal Images were taken on a Zeiss LSM 510 and processed using
243 ImageJ software. Images were taken of eight cells/groups of cells and the images presented in the
244 manuscript are representative of the sampled cell population.”

245

Results

BSO treatment

While maximising the desired effect of the BSO on the cells, it is important that the viability of the cells is not affected. TTV showed a 40% reduction in intracellular levels of GSH for the HaCaT cells upon 200µM BSO exposure for 18 hours (adapted from the methods of: He *et al.* (2003)).²⁴ This dose and time point were found to have a minimal impact on cellular viability as measured by the AB and MTT assays (data available in Supplementary Material, Figure: S3). Higher concentrations were found to have an effect on cellular viability, although a more pronounced effect was observed by CLSM, where signs of cellular stress were noted (data not shown). Therefore 200µM BSO exposure was chosen as optimal.

PAMAM G4 dendrimers

CLSM was employed as a way to examine the effect of BSO treatment on the cellular uptake mechanisms and subsequent oxidative stress. Early endosome formation was tracked with the Celllight® Early Endosome – RFP kit and the formation of ROS was also tracked using the carboxy-H₂DCFDA dye. Fluorescently labelled PSNP-NH₂ of 100nm diameter were employed as positive controls, and the results for those are shown in the Supplementary Material: Figure S2.




Figure 1: CLSM images of HaCaT (live) cells, upon exposure to 3.21 μ M PAMAM G4 dendrimer at 3 hours (no BSO is present in this sample). Image a) shows the bright field image of the cells, Image b) shows the fluorescence generated by the early endosomal red fluorescent protein, Image c) shows the fluorescence generated by the ROS (interacting with the carboxy- H_2 DCFDA dye) and Image d) shows the overlay of images a-c, where yellow coloured areas indicate simultaneous endosomal and ROS activity.

Figure 1 shows the HaCaT cells following the reported paradigm of PAMAM G4 endocytosis and subsequent ROS generation at the endosomal sites.^{6,7,8,9} Co-localisation (performed on the images in Figure 1) shows that 91(\pm 3)% of the generated ROS occurs in the neighbourhood of endosomes, and that 71(\pm 4)% of endosomal formation resulted in increases in ROS production (the other 30% of endosomal activity is most likely due to endocytosis which would be routinely done by the cell and would not involve the dendrimers and therefore not produce ROS).




Figure 2: Confocal images of HaCaT (live) cells, upon exposure to 3.21 μ M PAMAM G4 dendrimer at 3 hours, with pre-treatment of BSO: 200 μ M for 18hours. Image a) shows the bright field image of the cells, Image b) shows the fluorescence generated by the early endosomal red fluorescent protein, Image c) shows the fluorescence generated by the ROS (interacting with the carboxy-H₂DCFDA dye) and Image d) shows the overlay of images a-c, where yellow coloured areas indicate simultaneous endosomal and ROS activity.

In contrast, Figure 2 shows HaCaT cells which have been treated with BSO for 18 hours prior to PAMAM G4 exposure (BSO was still present in the media upon exposure, to ensure conditions were consistent). In images (b) and (c), the intensity of the endosomal RFP and ROS dye have been significantly reduced. Again co-localisation analysis was performed (on images in Figure 2) and showed that 30(\pm 6)% of the generated ROS was happening around the endosomes, and that 46(\pm 14)% of endosomal formation resulted in increases in ROS production. Intensity analysis of the red fluorescent protein produced by endocytosis, on average, showed a reduction in intensity of 70(\pm 3)% for cells treated with BSO.

Endosomal uptake was clearly reduced, although a decrease in the intensity of response of the ROS detection dye, carboxy-H₂DCFDA, was also noted, prompting a quantitative analysis of the ROS contents of the HaCaT cells: Figure 3.




Figure 3: Dose dependant generation of ROS in HaCaT cells (shown for 24 hours for the PAMAM G4). ROS is compared to BSO control which was set to equal 0%. The X-axis is plotted logarithmically to allow for better visualisation of lower concentrations. Data points are the mean of 18 samples, with error bars showing (\pm) the standard deviation.

The analysis of ROS showed a net decrease at all doses tested; for clarity, only the dose dependence of the 24 hour test has been shown, the full data-set is available in the supplementary material section (Figure S4). Over the dose range tested, the ROS response followed a monotonically decreasing trend. This result, is again, in contrast with previous observations, in which ROS levels were seen to increase above control levels in a dose dependant fashion, upon G4 dendrimer exposure.⁸

To examine how these changes in both endocytosis and ROS production affected the viability of the cell AB and MTT assays were performed. HaCaT cells were exposed to 200 μ M BSO for 18 hours and subsequently exposed to varying concentrations of PAMAM G4 nanoparticles. Viability was measured at 6, 12, 24, 48 and 72 hours (Figure 4 (a) and (b)).




Figure 4: Alamar Blue (a) and MTT (b) dose dependant viability results for PAMAM G4 dendrimers in HaCaT cells after 6, 12, 24, 48 and 72 hours. Viability is calculated as the percentage of living cells as compared to BSO control. Data points are the mean of 18 samples, with error bars showing (\pm) the standard deviation.

The AB assay shows little or no toxicity of PAMAM G4 at time points 6, 12, and 24 hours, whereas the viability is reduced to 50% at 48 hours and 30% at 72 hours, for the higher dose exposures. In HaCaT cells which have not been treated with BSO, the EC_{50} obtained from AB is around 10 μ M at 24 hours,⁷ whereas, in Figure 4, at the same time point, it is clear that there is no significant toxicity. The MTT results show a dramatically different dose dependent cytotoxicity profile for the BSO dosed cells compared to un-dosed.⁷ At 6 hours exposure (Figure 4b), the

viability is seen to be reduced to around 80% over the entire dose range. However, at exposure times of 12, 24, 48 and 72 hours, increases in viability are seen for the low-medium dose range. The effect is particularly pronounced at the 48 hour time point, at which the recorded MTT response is ~175% of control, for an exposure dose of ~1 μ M. For doses of ~2.6 μ M and higher, the MTT assay response registers a decrease in viability, which is more pronounced in the longer time point exposures.

PAMAM G6 dendrimers

Analysis of uptake and ROS production was again carried out using CLSM (with the same method as the PAMAM G4) incorporating the carboxy-H₂DCFDA dye and Celllight[®] Early Endosome – RFP kit (Figure 5 and 6).




Figure 5: Confocal images of HaCaT (live) cells, upon exposure to 1 μ M PAMAM G6 dendrimer at 1 hour (no BSO is present in this sample). Image a) shows the bright field image of the cells, Image b) shows the fluorescence generated by the early endosomal red fluorescent protein, Image c) shows the fluorescence generated by the ROS (interacting with the carboxy- H_2 DCFDA dye) and Image d) shows the overlay of images a-c, where yellow coloured areas indicate simultaneous endosomal and ROS activity.

In the absence of BSO, PAMAM G6 exposure, resulted in strong red fluorescence indicating a high level of endocytosis, complemented by strong green fluorescence indicating increased ROS production. Co-localisation analysis (of images in Figure 5) found that 75(\pm 2)% of the generated ROS occurred in the region of the endosomes, and that 92(\pm 1)% of endocytosis resulted in increases in ROS production. This strong level of co-localisation is indicative of the accepted paradigm of nanoparticle uptake by endocytosis, and ROS production at the site of endosomes.




Figure 6: Confocal images of HaCaT (live) cells, upon exposure to 1 μ M PAMAM G6 dendrimer at 1 hour, with pre-treatment of BSO: 200 μ M. Image a) shows the bright field image of the cells, Image b) shows the fluorescence generated by the early endosomal red fluorescent protein, Image c) shows the fluorescence generated by the ROS (interacting with the carboxy-H₂DCFDA dye) and Image d) shows the overlay of images a-c, where yellow coloured areas indicate simultaneous endosomal and ROS activity.

In comparison to exposure of PAMAM G6 in cells without BSO treatment (untreated cells), there is a sparsity of endosomal activity and oxidative stress response in cells which had been pre-treated with BSO for 18 hours. Analysis (of the images in Figure 6) showed decreased levels of co-localisation between the ROS and endosomal formation: only 41(\pm 18)% of ROS production was recorded in areas where endosomes had formed and 55(\pm 10)% of endocytosis resulted in increases in ROS production. Intensity analysis of the red fluorescent protein produced by endocytosis, on average, showed a reduction in intensity of 61(\pm 1)% for cells treated with BSO when compared to the untreated cells.

Again, in light of the reduced ROS response, quantitative analysis, using carboxy-H₂DCFDA was performed: Figure 7.




Figure 7: Dose dependant generation of ROS in HaCaT cells (shown at 24 hours for the PAMAM G6). ROS is compared to BSO control which was set to equal 0%. The X-axis is plotted logarithmically to allow for better visualisation of lower concentrations. Data points are the mean of 18 samples, with error bars showing (\pm) the standard deviation.

The dose response of the ROS generation shows similarities with the G4 in the overall progression of the response, pointing to a trend where increasing concentration leads to decreasing amounts of ROS, contrary to what has been observed in previous studies without BSO exposure.⁸ As with the G4 analysis, for clarity, only one of the eight concentrations tested have been shown, the full data-set is available in the supplementary material section (Figure S5). At all doses tested, the ROS response followed the same trend as shown in Figure 7.

To analyse the effect this had on viability, AB and MTT assays were performed: Figure 8.




Figure 8: Alamar Blue (a) and MTT (b) dose dependent toxicity results for PAMAM G6

dendrimers in HaCaT cells after 6, 12, 24, 48 and 72 hours. Viability is calculated as the

percentage of living cells as compared to a BSO control. The X-axis is plotted logarithmically to

allow for better visualisation of lower doses. Data points are the mean of 18 samples, with error

bars showing (\pm) the standard deviation.

The viability results of the PAMAM G6 dendrimers are similar to those observed in the G4

analysis. In Figure 8, for 6 and 12 hour exposure, little toxicity is recorded by AB, although the

response is significantly stronger at the later time points. The previously reported EC_{50} obtained

from the AB assay for PAMAM G6 dendrimers is in the range of 1 μM-1.6 μM at 24 hours,⁸

which would appear to agree with the values obtained in this study. However, as was the case for

PAMAM G4 exposure, the MTT assay registers increased percentage viability, compared to

BSO control, for the intermediate doses, for all but the 6 hour exposure time point. A cytotoxic

response is elicited for doses greater than 1 μM , for the 24, 48 and 72 hour time points and the response is consistent with the previously reported EC_{50} values of 0.92 μM -1.13 μM .⁷

PAMAM G4-G6 Comparative Analysis

PAMAM nanoparticle toxicity is a generation dependant process,^{7,8,9} therefore, a comparison of the response of BSO treated cells to G4 and G6 dendrimer exposure was performed, to establish whether a similar generation dependence of the cellular responses is still evident.

The AB and MTT assays were compared for PAMAM G4 and G6 dendrimers, and the results are shown in Figures 9 (AB) and 10 (MTT). For both dendrimers, approximately equivalent concentration ranges were used, although, for the PAMAM G6 dendrimers, a slightly lower initial concentration was used due to the higher associated toxicity.^{7,8,9,10}




Figure 9: Alamar Blue (AB) dose dependant toxicity results comparing the PAMAM G4 and G6 dendrimers in HaCaT cells after 48 hours (a) and 72 hours (b). Viability is calculated as the percentage of living cells as compared to BSO control. The x-axis is plotted logarithmically to

allow for better visualisation of lower doses. Data points are the mean of 18 samples, with error bars showing (\pm) the standard deviation. 6, 12 and 24 hour graphs can be seen in the supplementary material section (Figure S6).

Little or no significant cytotoxicity was registered by the AB assay for either dendrimer at the 6 and 12 hour time points. At 24 hours, the higher toxicity associated with the G6 dendrimers becomes apparent, while the G4 still shows no significant change (Supplementary Material: Figure S6, c). At 48 hours (Figure 9(a)) the G4 dendrimers begin to elicit a significant toxic response and finally, at 72 hours (Figure 9(b)), the G4 toxic profile is beginning to match that of G6, although the reduction of viability is much higher for G6.




Figure 10: MTT dose dependant toxicity results comparing the PAMAM G4 and G6 dendrimers in HaCaT cells after 48 hours (a) and 72 hours (b). Viability is calculated as the percentage of living cells as compared to BSO control. The X-axis is plotted logarithmically to allow for better visualisation of lower doses. Data points are the mean of 18 samples, with error bars showing (\pm)

the standard deviation. 6, 12 and 24 hour graphs can be seen in the supplementary material section (Figure S7).

In the MTT, similar to AB response, no significant difference was seen in the 6 hour analysis for the two dendrimers, although differences become apparent at 12 hours (Supplementary Material: Figure S7, b), at which point the MTT response increases above that of controls for the G6 dendrimer. This increase in MTT response also occurs for the G4 dendrimers, although, it is not manifest until the later time point of 48 hours (Figure 10(a)). Similar to the AB response, cytotoxicity is registered by the MTT assay for both dendrimers beginning at 24 hours; however, the AB only reaches about 50% viability. At 48 and 72 hours a more complete toxic profile for both assays is seen and distinct generation dependence is observable, more consistent with that observed for untreated cells.⁷




Figure 11: Dose dependant generation of ROS in HaCaT cells (shown at 24 hours for PAMAM G4 and G6). Data is compared to BSO control which was set to equal 0%. The X-axis is plotted logarithmically to allow for better visualisation of lower concentrations. Data points are the mean of 18 samples, with error bars showing (\pm) the standard deviation.

To further investigate any generation dependence, the dose response of the ROS was compared for the G4 and the G6 (Figure 11). It is notable that the generation dependence of the trends is reversed in Figure11 where G4 elicits a more pronounced reduction in ROS levels than G6.

Discussion

The confocal images of Figures: 1 and 2 (PAMAM G4) and Figures: 5 and 6 (PAMAM G6) show that, for cells treated with 200 μ M BSO for 18 hours prior to PAMAM exposure, the rate of endocytosis has been markedly reduced. This reduction in endocytosis is accompanied by a similar reduction in intracellular ROS and a dramatic change in the responses of the cytotoxic assays. Nevertheless, the observed reduction of intracellular ROS and cytotoxic responses are systematically dependent on dendrimer exposure time, dose and generation, consistent with the intracellular action of the dendritic nanoparticles.

The demonstrated mechanism of PAMAM dendrimer toxicity is one of endocytosis,^{8,43,44} ROS production,^{7,8,9,18} subsequent endosomolysis, whereby the nanoparticle bursts out of the endosome/lysosome into the cytosol,¹⁹ and localisation in the mitochondria.⁴⁵ Mukherjee and Byrne (2013) identified two apoptotic pathways, the death-receptor pathway (extrinsic, Fas mediated FADD pathway⁴⁶) and the mitochondrial pathway (intrinsic, TNF- α mediated FADD⁴⁷).¹⁰ The former is initiated by the earlier stage ROS generation in the region of the endosomes, while the latter is initiated by the localisation of the dendrimers in the mitochondria. It has been proposed that the early stage ROS production is due to the action of NADPH oxidase⁶ (producing superoxide anions (O_2^-)⁴⁸) and the v-ATPase proton pump (providing protons,⁴⁹ ultimately leading to production of H_2O_2 ⁵⁰ in and around the endosome. In cells treated with BSO, however, PAMAM dendrimers elicit a dramatically different cytotoxicity profile, as registered by the AB and MTT assays, compared to that of untreated cells.^{7,8}

Studies by Khalid *et al.* (2015) of cellular uptake of PPI dendrimers have demonstrated that, although the larger generation PPI dendrimers are endocytosed and elicit similar responses to PAMAM equivalents in HaCaT cells, for smaller generation PPI dendrimers, uptake by passive diffusion occurs and, when the dendrimers enter the cell in this way, they were observed to act as antioxidants and elicit a significantly reduced cytotoxic effect.¹⁸ BSO exposure has been shown to permeabilise the cell membrane,²⁸ and as a result, it is proposed that the PAMAM G4 and G6 dendrimers are able to circumvent the endocytotic process, are uptaken by passive diffusion, and, as PAMAM dendrimers are similar in structure to PPI dendrimers and have comparable surface chemistry, similarly behave as antioxidants in the cytosol, eliciting substantially reduced cytotoxic responses.

Alamar Blue is a non-specific assay and measures cellular viability based on the overall activity of the cytosolic environment.⁵¹ The significant reduction, rather than expected increase, of toxicity as registered by this assay reflects the reduction in the endocytosis process (which initiates the Fas mediated FADD (death-receptor) apoptotic pathway), in favour of the passive diffusion of nanoparticles across the membrane, and consequent reduction in ROS generation in the region of endosomes. However, endocytosis is not fully eliminated for either generation, and therefore, the activation of the Fas mediated pathway, on a much reduced scale, would in part explain why the generation dependent response is still observed for the PAMAMs.

The MTT assay measures mitochondrial activity as an indicator of cellular viability^{52,53,54} and, in the case of the studies described here, the mitochondria are implicated in at least two different processes and changes in MTT responses reflect the dose and generation dependence of these

processes at several time points. The first process is the loss of GSH from the cell, which has been shown to cause activation of mitochondrial signalling pathways and expression of genes associated with apoptosis, growth and differentiation.^{33,38,39,40} This overall increase in mitochondrial activity (observed in the low dose regime), is seen as the initial increase in MTT values above controls, associated with dose and generation dependent decrease in ROS below controls, due to the antioxidant effect of the passively uptaken dendrimer nanoparticles.

The second effect gives rise to a sharp decrease in mitochondrial activity (observed in the higher dose regime); consistent with PAMAM dendrimer localisation and disruption of the mitochondria,^{7,8} initiating the mitochondrial apoptotic pathway, leading to cell death in a dose dependant fashion, as observed for untreated cells.^{7,8} This process may be accelerated via the opening of the mitochondrial membrane permeability transition pore.^{30,31,32} Whether passively diffused into the cell, or released into the cytosol by endosomolysis after endocytosis, the result of free PAMAM dendrimers in the cytosol and subsequent localisation to the mitochondria should be equivalent, both resulting in disruption of the mitochondria, a second phase increase in ROS within the cell, subsequent decay in the mitochondrial membrane potential and finally the initiation of a cascade leading to apoptosis.^{7,8,9,10} It should also be noted that in both cases the opening of the mitochondrial membrane transition pore occurs, either by the action of BSO,^{30,31,32} or due to the release of the endosomal/lysosomal contents causing intracellular release of Ca^{2+} , leading to calcium dependant opening of the pore.²¹ This would, in both cases, facilitate the entry of the dendrimer to the mitochondria. As a result, in the high dose regime, the observed toxic response of the BSO treated cells, as registered by the MTT assay, is not very different to that observed for untreated cells.

534
535 The passive diffusion of dendrimers across the cell membrane is a size dependant process¹⁸ and,
536 the generation dependence of the cellular responses to the two PAMAM dendrimer generations,
537 G4 and G6, (Figures: 1, 2, 5 and 6), is consistent with a higher uptake rate for the G4 dendrimer
538 than the G6. The greater reduction of intensity of Rab-5a-RFP compared to controls (with no
539 BSO treatment), G4 dendrimers (70%) compared to G6 (60%), indicates a higher diffusion rate
540 for G4 dendrimers, leaving fewer available for endocytosis. This further explains the higher rate
541 of anti-oxidative activity exhibited by the G4 dendrimers when compared to G6 (Figure 11).
542 Overall, it would appear that the membrane has become more permeable to an extent that favours
543 passive uptake, although not completely eliminating active endocytosis.

544
545 Within the framework of Adverse Outcome Pathways (AOPs), recently endorsed by the
546 OECD^{55,56} as a method to simplify the representation of the mode of action of a toxicant or
547 agonist, the generation of ROS can be considered the key Molecular Initiating Event (MIE) of
548 the AOP, which ultimately leads to the AO of loss of cell viability. The treatment with BSO
549 causes a depletion of GSH, which would lead to the expectation of much increased ROS levels
550 after endocytosis. However, that was not observed and the result of the reduction of intrinsic
551 GSH levels by BSO treatment was predominantly the increased permeability of the cell
552 membrane, resulting in an increased rate of uptake of the PAMAM dendrimers by passive
553 diffusion, making it a favoured uptake mechanism, more so for the smaller G4 dendrimer than
554 the larger G6. The co-existence of the parallel uptake mechanisms increases the complexity of
555 any model to describe the *in vitro* system, although it could prove a useful model to develop
556 networks of AOPs, initiated by different MIEs. However, to fully examine the effect of reduction

of GSH in terms of decreased anti-oxidant activity alone, an assay which did not cause decrease of the permeabilisation of the cell membrane would be necessary.

Considering the potential for PAMAM dendrimers in nanomedical applications, it is important to note that, when diffused into the cell, the aminated surface chemistry of the dendrimers lend them antioxidant activity, similar to small molecular anti-oxidants, such as N-acetylcysteine (NAC) and NAC amide (NACA - a more bio-available version of NAC). NACA has been extensively studied as an antioxidant in the cell, due to its ability to diffuse across the membrane and the presence of a terminal proton donor group.^{57,58} Interestingly, it has also been shown to completely reverse the damage caused to the cell by depletion of GSH.^{59,60,61} NACA, due to this strong anti-oxidant ability, has been proposed in the treatment of several disorders and diseases, such as: HIV,⁶² Alzheimer's and Parkinson's disease,^{60,63} cataract formation,⁵⁹ retinal degeneration⁶⁴ and essentially any disease where ROS is identified as the potential MIE.^(See: 61 and references therein) If PAMAM dendrimers are seen to act in a similar way, it may potentially allow for their use in a plethora of different nano-medical applications. As a strategy for drug release, endosomolysis can be extremely disruptive to the cell²⁰ and therefore, in the case of cationic nanoparticles for intracellular nanomedical applications, avoiding the process of endocytosis may be a valid strategy to pursue.²² In terms of therapeutic applications, direct entry into the cytosol may be a more convenient route for drug or gene delivery.

577 Conclusions

578 Although PAMAM dendritic nanoparticles are known to elicit significant cytotoxic responses *in*
579 *vitro*, the cellular response mechanisms can be notably altered by treatment of the cells with
580 BSO. The treatment increases the cell membrane permeability, enabling uptake of the particles
581 by passive diffusion, where after, they act as antioxidants in the cytosol, rather than producing
582 oxidative stress in the region of endosomes. The ability to tune the cellular uptake mechanism
583 allows direct entry into the cytosol and may have important implications for nanotoxicity as well
584 as drug and gene delivery using nanovehicles.

585

586 Acknowledgements

587 The authors would like to thank Dr. Alan Casey and Esen Efeoglu for their contributions in the
588 capturing and processing of the confocal images for this manuscript.

589

590 Funding Sources

591 Irish Government's Programme for Research in Third Level Institutions, Cycle 4, National
592 Development Plan 2007-2013, supported by the European Union Structural Fund. M Maher is
593 funded through the DIT Fiosraigh President's Award for Research Excellence 2010

594

595 Declaration of interest

596 The authors declare no conflict of interest related to the work presented in this manuscript.

597

References

- 1: Eichman, J.D., Bielinska, A.U., Kukowska-Latallo, J.F. and Baker, J.R.Jr. (2000) The use of PAMAM dendrimers in the efficient transfer of genetic material into cells. *J. Pharm. Sci. Technol.* 3(7): 233-245.
- 2: Duncan, R. and Izzo, L. (2005) Dendrimer biocompatibility and toxicity. *Adv. Drug. Deliv. Rev.* 57(15): 2215– 2237.
- 3: De Jong, W. H. and Borm, P.J.A. (2008) Drug delivery and nanoparticles: Applications and hazards. *Int. J. Nanomedicine.* 3(2): 133–149.
- 4: Nel, A., Xia, T., Mädler, L. and Li, N. (2006) Toxic potential of materials at the nanolevel. *Science.* 311(5761): 622-627.
- 5: Jain, K.K., Kesharwani, P., Gupta, U. and Jain, N.K. (2010) Dendrimer toxicity: Let's meet the challenge. *Int. J. Pharm.* 394(1-2): 122–142.
- 6: Xia, T., Kovochich, M., Brant, J., Hotze, M., Sempf, J., Oberley, T., Sioutas, C., Yeh, J.I., Wiesner, M.R. and Nel, A.E. (2006) Comparison of the Abilities of Ambient and Manufactured Nanoparticles to Induce Cellular Toxicity According to an Oxidative Stress Paradigm. *Nano Lett.* 6(8): 1794-1807.

620 7: Mukherjee, S.P., Davoren, M. and Byrne, H.J. (2010) *In vitro* mammalian cytotoxicological
621 study of PAMAM dendrimers - towards quantitative structure activity relationships. *Toxicol. in*
622 *Vitro*, 24(1): 169-177.

623

624 8: Mukherjee, S.P., Lyng, F.M., Garcia, A., Davoren, M. and Byrne H.J. (2010) Mechanistic
625 studies of *in vitro* cytotoxicity of Poly(amidoamine) dendrimers in mammalian cells. *Toxicol.*
626 *Appl. Pharmacol.* 248(3): 259–268.

627

628 9: Naha, P.C., Davoren, M., Lyng, F.M. and Byrne, H.J. (2010) Reactive Oxygen Species (ROS)
629 induced cytokine production and cytotoxicity of PAMAM dendrimers in J774A.1 cells. *Toxicol.*
630 *Appl. Pharmacol.* 246(1-2): 91-99

631

632 10: Mukherjee S.P. and Byrne, H.J. (2013) Polyamidoamine Dendrimer Nanoparticle
633 Cytotoxicity, Oxidative Stress, Caspase Activation and Inflammatory Response: Experimental
634 Observation and Numerical Simulation. *Nanomed. Nanotech. Biol. Med.* 9(2): 202-211.

635

636 11: Anguissola, S., Garry, D., Salvati, A., O'Brien, P.J. and Dawson, K.A. (2014) High Content
637 Analysis Provides Mechanistic Insights on the Pathways of Toxicity Induced by Amine-
638 Modified Polystyrene Nanoparticles. *PLoS ONE*, 9(9): e108025.
639 doi:10.1371/journal.pone.0108025

640

641 12: Paget, V., Dekali, S., Kortulewski, T., Grall, R., Gamez, C., Blazy, K., Aguerre-Chariol, O.,
642 Chevillard, S., Braun, A., Rat, P and Lacroix, G. (2015) Specific Uptake and Genotoxicity

643 Induced by Polystyrene Nanobeads with Distinct Surface Chemistry on Human Lung Epithelial
 644 Cells and Macrophages. *PLoS ONE*, 10(4): e0123297. doi:10.1371/journal.pone.0123297
 645
 646 13: Lunov, O., Syrovets, T., Loos, C., Beil, J., Delacher, M., Tron, K., Nienhaus, G.U.,
 647 Musyanovych, A., Mailänder, V., Landfester, K. and Simmet, T. (2011) Differential Uptake of
 648 Functionalized Polystyrene Nanoparticles by Human Macrophages and a Monocytic Cell Line.
 649 *ACS Nano*. 5(3): 1657-1669.
 650
 651 14: Nabeshi, H., Yoshikawa, T., Matsuyama, K., Nakazato, Y., Tochigi, S., Kondoh, S., Hirai,
 652 T., Akase, T., Nagano, K., Abe, Y., Yoshioka, Y., Kamada, H., Itoh, N., Tsunoda, S. and
 653 Tsutsumi, Y. (2011) Amorphous nanosilica induce endocytosis dependent ROS generation and
 654 DNA damage in human keratinocytes. *Part. Fibre Toxicol.* 8(1): 10 pages.
 655
 656 15: Mozdzan, M., Szemraj, J., Rysz, J., Stolarek, R.A. and Nowak, D. (2006) Anti-oxidant
 657 activity of spermine and spermidine re-evaluated with oxidising system involving iron and
 658 copper ions. *Int. J. Biochem. Cell Biol.* 38(1): 69-81.
 659
 660 16: Lotito, S.B. and Frei, B. (2004) Relevance of apple polyphenols as antioxidants in human
 661 plasma: contrasting in vitro and in vivo effects. *Free Radic. Biol. Med.* 36(2): 201-211.
 662
 663 17: Hipkiss AR. (2009) Carnosine and its possible roles in nutrition and health. *Adv. Food Nutr.*
 664 *Res.* 57(-): 87-154.
 665

666 18: Khalid, H., Mukherjee, S.P., O'Neill, L. and Byrne, H.J. (2015) Structural dependence of the
 667 *In vitro* cytotoxicity, oxidative stress and uptake mechanisms of Poly(propylene imine) dendritic
 668 nanoparticles. *J. Appl. Toxicol.* (in press - Accepted: October 2015)
 669
 670 19: Watson, P., Jones, A.T. and Stephens, D.J. (2005) Intracellular trafficking pathways and drug
 671 delivery: fluorescence imaging of living and fixed cells. *Adv. Drug Deliver. Rev.* 57(1): 43-61.
 672
 673 20: Mukherjee, S.P. (2012) Towards Structure Activity Relationships for *in vitro* Toxicity of
 674 Polyamidoamine Dendritic Nanoparticles. *Thesis (PhD)*, Dublin Institute of Technology: 2012.
 675 Dublin: DIT.
 676
 677 21: Xia, T., Kovochich, M., Liong, M., Mädler, L., Gilbert, B., Shi, H., Yeh, J.I., Zink, J.I. and
 678 Nel, A.E. (2008) Comparison of the mechanism of toxicity of zinc oxide and cerium oxide
 679 nanoparticles based on dissolution and oxidative stress properties. *ACS Nano.* 2(10): 2121-2134.
 680
 681 22: Guarnieri, D., Sabella, S., Muscetti, O., Belli, V., Malvindi, M.A., Fusco, S., De Luca, E.,
 682 Pompa, P.P. and Netti, P.A. (2014) Transport across the cell-membrane dictates nanoparticle fate
 683 and toxicity: a new paradigm in nanotoxicology. *Nanoscale.* 6(17): 10264-10273.
 684
 685 23: Girard, P-M., Graindorge, D., Smirnova, V., Rigolet, P., Francesconi, S., Scanlon, S. and
 686 Sage, E. (2013) Oxidative Stress in Mammalian Cells Impinges on the Cysteines Redox State of
 687 Human XRCC3 Protein and on Its Cellular Localization. *PLoS ONE*, DOI:
 688 10.1371/journal.pone.0075751

689

690 24: He, Y-Y., Huang, J-L., Ramirez, D.C. and Chignell, C.F. (2003) Role of Reduced
691 Glutathione Efflux in Apoptosis of Immortalized Human Keratinocytes Induced by UVA. *J.*
692 *Biol. Chem.* 278(10): 8058–8064.

693

694 25: Madesh, M., Benard, O. and Balasubramanian, K.A. (1998) Glutathione modulates lipid
695 composition of human colon derived HT-29 cells. *Int. J. Biochem. Cell Biol.* 30(12): 1345-1352.

696

697 26: Tobi, S.E., Paul, N. and McMillan T.J. (2000) Glutathione modulates the level of free
698 radicals produced in UVA-irradiated cells. *J. Photochem. Photobiol. B.* 57(2-3): 102–112.

699

700 27: Li, X.Y., Donaldson, K., Rahman, I. and Mac Nee, W. (1994) An investigation of the role of
701 glutathione in increased epithelial permeability induced by cigarette smoke *in vivo* and *in vitro*.
702 *Am. J. Respir. Crit. Care Med.* 149(6): 1518-1525.

703

704 28: Zaman, G.J., Lankelma, J., van Tellingen, O., Beijnen, J., Dekker, H., Paulusma, C., Oude
705 Elferink, R.P., Baas, F. and Borst, P. (1995) Role of glutathione in the export of compounds from
706 cells by the multidrug-resistance-associated protein. *Proc. Natl. Acad. Sci. USA.* 92(17): 7690-
707 7694.

708

709 29: Griffith, O.W. (1982) Mechanism of Action, Metabolism, and Toxicity of Buthionine
710 Sulfoximine and Its Higher Homologs, Potent Inhibitors of Glutathione Synthesis. *J. Biol. Chem.*
711 257(22): 13704-13712.

712

713 30: Armstrong, J.S. and Jones, D.P. (2002) Glutathione depletion enforces the mitochondrial
714 permeability transition and causes cell death in Bcl-2 overexpressing HL60 cells. *FASEB J.*
715 *16(10)*: 1263-1265.

716

717 31: Chernyak, B.V. and Bernardi, P. (1996) The mitochondrial permeability transition pore is
718 modulated by oxidative agents through both pyridine nucleotides and glutathione at two separate
719 sites. *Eur. J. Biochem.* *238(3)*: 623–630.

720

721 32: Reed, D.J. and Savage, M.K. (1995) Influence of metabolic inhibitors on mitochondrial
722 permeability transition and glutathione status. *Biochim. Biophys. Acta.* *1271(1)*: 43–50.

723

724 33: Circu, M.L. and Aw, T.Y. (2008) Glutathione and apoptosis. *Free Radic. Res.* *42(8)*: 689–
725 706.

726

727 34: Lash, L.H. (2006) Mitochondrial Glutathione Transport: Physiological, Pathological and
728 Toxicological Implications. *Chem. Biol. Interact.* *163(1-2)*: 54–67.

729

730 35: Beatrice, M.C., Stiers, D.L. and Pfeiffer, D.R. (1984) The role of glutathione in the retention
731 of Ca^{2+} by liver mitochondria. *J. Biol. Chem.* *259(2)*: 1279–1287.

732

733 36: Lotscher, H.R., Winterhalter, K.H., Carafoli, E. and Richter, C. (1979) Hydroperoxides can
734 modulate the redox state of pyridine nucleotides and the calcium balance in rat liver
735 mitochondria. *Proc. Natl. Acad. Sci. USA*. 76(9): 4340–4344.
736

737 37: Olafsdottir, K., Pascoe, G.A. and Reed, D.J. (1988) Mitochondrial glutathione status during
738 Ca²⁺ ionophore-induced injury to isolated hepatocytes. *Arch. Biochem. Biophys.* 263(1): 226–
739 235.
740

741 38: Davis, W.Jr., Ronai, Z. and Tew, K.D. (2001) Cellular thiols and reactive oxygen species in
742 drug-induced apoptosis. *J. Pharmacol. Exp. Ther.* 296(1): 1–6.
743

744 39: Petit, P.X., Susin, S.A., Zamzami, N., Mignotte, B. and Kroemer, G. (1996) Mitochondria
745 and programmed cell death: Back to the future. *FEBS Lett.* 396(1): 7–13.
746

747 40: Skulachev, V.P. (1996) Why are mitochondria involved in apoptosis? Permeability transition
748 pores and apoptosis as selective mechanisms to eliminate superoxide-producing mitochondria
749 and cells. *FEBS Lett.* 397(1): 7–10.
750

751 41: Maher, M.A., Naha, P.C., Mukerjee, S.P. and Byrne, H.J. (2014) Numerical simulations of in
752 vitro nanoparticle toxicity – the case of Poly(amido amine) dendrimers. *Toxicol. In Vitro.* 28(8):
753 1449-1460.
754

755 42: Zinchuk, V. and Zinchuk, O. (2008) Quantitative colocalization analysis of confocal
756 fluorescence microscopy images. *Curr. Protoc. Cell Biol. Ch.4(Unit 4.19)*, 1-16.
757

758 43: Kitchens, K.M., Foraker, A.B., Kolhatkar, R.B., Swaan, P.W., Ghandehari, H. (2007)
759 Endocytosis and interaction of poly (Amidoamine) dendrimers with Caco-2 cells. *Pharmaceut.*
760 *Res. 24(11)*: 2138–2145.
761

762 44: Hong, S., Bielinska, A.U., Mecke, A., Keszler, B., Beals, J.L., Shi, X., Balogh, L., Orr, B.G.,
763 Baker, J.R.Jr and Banaszak Holl, M.M. (2004) Interaction of poly(amidoamine) dendrimers with
764 supported lipid bilayers and cells: hole formation and the relation to transport. *Bioconjugate*
765 *Chem. 15(4)*: 774–782.
766

767 45: Lee, J.H., Cha, K.E., Kim, M.S., Hong, H.W., Chung, D.J., Ryu, G. and Myung, H. (2009)
768 Nanosized polyamidoamine (PAMAM) dendrimer-induced apoptosis mediated by mitochondrial
769 dysfunction. *Toxicol. Lett. 190(2)*: 202–207.
770

771 46: Wajant, H. (2002) The Fas signaling pathway: more than a paradigm. *Science*
772 *296(5573)*:1635-1636.
773

774 47: Desagher, S. and Martinou, J.C. (2000) Mitochondria as the central control point of
775 apoptosis. *Trends Cell Biol. 10(9)*: 369-377.
776

777 48: Shuvaev, V.V., Han, J., Yu, K.J., Huang, S., Hawkins, B.J., Madesh, M., Nakada, M. and
778 Muzykantov, V.R. (2011) PECAM-targeted delivery of SOD inhibits endothelial inflammatory
779 response. *FASEB J.* 25(1): 348-357.
780

781 49: Nel, A.E., Mädler, L., Velegol, D., Xia, T., Hoek, E.M.V., Somasundaran, P., Klaessig, F.,
782 Castranova, V. and Thompson, M. (2009) Understanding biophysicochemical interactions at the
783 nano–bio interface. *Nature Materials.* 8(7): 543 – 557.
784

785 50: Babior, B.M., Lambeth, J.D. and Nauseef, W. (2002) The neutrophil NADPH oxidase. *Arch.*
786 *Biochem. Biophys.* 397(2): 342–344.
787

788 51: O'Brien, J., Wilson, I., Orton, T. and Pognan, F. (2000) Investigation of the Alamar Blue
789 (resazurin) fluorescent dye for the assessment of mammalian cell cytotoxicity. *Eur. J. Biochem.*
790 267(17): 5421-5426.
791

792 52: Mosmann, T. (1983) Rapid colorimetric assay for cellular growth and survival: Application
793 to proliferation and cytotoxicity assays. *J. Immunol. Methods.* 65(1): 55-63.
794

795 53: Denizot, F. and Lang, R. (1986) Rapid colorimetric assay for cell growth and survival:
796 Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. *J.*
797 *Immunol. Methods.* 89(2): 271-277.
798

799 54: Hansen, M.B., Nielsen, S.E. and Berg, K. (1989) Re-examination and further development of
800 a precise and rapid dye method for measuring cell growth/cell kill. *J. Immunol. Methods.* 119(2):
801 203-210.

802

803 55: Organisation for Economic Co-Operation and Development (OECD): Guidance Document
804 on the Validation of (Quantative) Structure Activity Relationships [(Q)SAR] Models, *OECD*,
805 *ENV/JM/MONO(2007)2*, 2007.

806

807 56: Organisation for Economic Co-Operation and Development (OECD): Guidance Document
808 On Developing and Assessing Adverse Outcome Pathways, *OECD*, *ENV/JM/MONO(2013)6*,
809 2013.

810

811 57: Price, T.O., Uras, F., Banks, W.A. and Ercal, N. (2006) A novel antioxidant N-acetylcysteine
812 amide prevents gp120- and Tat-induced oxidative stress in brain endothelial cells. *Exp. Neurol.*
813 *201(1)*: 193–202.

814

815 58: Grinberg, L., Fibach, E., Amer, J. and Atlas, D. (2005) N-acetylcysteine amide, a novel cell
816 permeating thiol, restores cellular glutathione and protects human red blood cells from oxidative
817 stress. *Free Radic. Biol. Med.* 38(1): 136–145.

818

819 59: Carey, J.W., Pinarci, E.Y., Penugonda, S., Karacal, H. and Ercal, N. (2011) *In vivo* inhibition
820 of l-buthionine-(S,R)-sulfoximine-induced cataracts by a novel antioxidant, N-acetylcysteine
821 amide. *Free Radic. Biol. Med.* 50(6): 722–729.

822

823 60: Penugonda, S., Mare, S., Goldstein, G., Banks, W.A. and Ercal, N. (2005) Effects of N-

824 acetylcysteine amide (NACA), a novel thiol antioxidant against glutamate-induced cytotoxicity

825 in neuronal cell line PC12. *Brain Research. 1056(2)*: 132–138.

826

827 61: Sunitha, K., Hemshekhar, M., Thushara, R.M., Santhosh, M.S., Yariswamy, M., Kemparaju,

828 K. and Girish, K.S. (2013) N-Acetylcysteine amide: a derivative to fulfill the promises of N-

829 Acetylcysteine. *Free Radic. Res. 47(5)*: 357-367.

830

831 62: Banerjee, A., Zhang, X., Manda, K.R., Banks, W.A. and Ercal, N. (2010) HIV proteins

832 (gp120 and Tat) and methamphetamine in oxidative stress-induced damage in the brain: potential

833 role of the thiol antioxidant N-acetylcysteine amide. *Free Radic. Biol. Med. 48(10)*: 1388-1398.

834

835 63: Lee, K.S., Kim, S.R., Park, H.S., Park, S.J., Min, K.H., Lee, K.Y., Choe, Y.H., Hong, S.H.,

836 Han, H.J., Lee, Y.R., Kim, J.S., Atlas, D. and Lee, Y.C. (2007) A novel thiol compound, N-

837 acetylcysteine amide, attenuates allergic airway disease by regulating activation of NF-kappaB

838 and hypoxia-inducible factor-1alpha. *Exp. Mol. Med. 39(6)*: 756-768.

839

840 64: Schimel, A.M., Abraham, L., Cox, D., Sene, A., Kraus, C., Dace, D.S., Ercal, N. and Apte,

841 R.S. (2011) N-acetylcysteine amide (NACA) prevents retinal degeneration by up-regulating

842 reduced glutathione production and reversing lipid peroxidation. *Am. J. Pathol. 178(5)*: 2032-

843 2043.